ON THE GROUP OF AUTOMORPHISMS OF THE BRANDT λ^0-EXTENSION OF A MONOID WITH ZERO

OLEG GUTIK

arXiv:1609.06085v1 [math.GR] 20 Sep 2016

Abstract. The group of automorphisms of the Brandt λ^0-extension $B^0_\lambda(S)$ of an arbitrary monoid S with zero is described. In particular we show that the group of automorphisms $\mathbf{Aut}(B^0_\lambda(S))$ of $B^0_\lambda(S)$ is isomorphic to a homomorphic image of the group defines on the Cartesian product $\mathscr{S}_\lambda \times \mathbf{Aut}(S) \times H_1^\lambda$ with the following binary operation:

$$[\varphi, h, u] \cdot [\varphi', h', u'] = [\varphi\varphi', hh', \varphi u' \cdot uh'],$$

where $\mathscr{S}_\lambda$ is the group of all bijections of the cardinal λ, $\mathbf{Aut}(S)$ is the group of all automorphisms of the semigroup S and H_1^λ is the direct λ-power of the group of units H_1 of the monoid S.

1. Introduction and preliminaries

Further we shall follow the terminology of [2, 21].

Given a semigroup S, we shall denote the set of idempotents of S by $E(S)$. A semigroup S with the adjoined unit (identity) [zero] will be denoted by S^1 [S^0] (cf. [2]). Next, we shall denote the unit (identity) and the zero of a semigroup S by 1_S and 0_S, respectively. Given a subset A of a semigroup S, we shall denote by $A^* = A \setminus \{0_S\}$.

If S is a semigroup, then we shall denote the subset of idempotents in S by $E(S)$. If $E(S)$ is closed under multiplication in S and we shall refer to $E(S)$ a *band* (or the *band of* S). If the band $E(S)$ is a non-empty subset of S, then the semigroup operation on S determines the following partial order $\leqslant$ on $E(S)$: $e \leqslant f$ if and only if $ef = fe = e$. This order is called the *natural partial order* on $E(S)$.

If $h\colon S \to T$ is a homomorphism (or a map) from a semigroup S into a semigroup T and if $s \in S$, then we denote the image of s under h by $(s)h$.

Let S be a semigroup with zero and λ a cardinal $\geqslant 1$. We define the semigroup operation on the set $B_\lambda(S) = (\lambda \times S \times \lambda) \cup \{0\}$ as follows:

$$(\alpha, a, \beta) \cdot (\gamma, b, \delta) = \begin{cases} (\alpha, ab, \delta), & \text{if } \beta = \gamma; \\ 0, & \text{if } \beta \neq \gamma, \end{cases}$$

and $(\alpha, a, \beta) \cdot 0 = 0 \cdot (\alpha, a, \beta) = 0 \cdot 0 = 0$, for all $\alpha, \beta, \gamma, \delta \in \lambda$ and $a, b \in S$. If $S = S^1$ then the semigroup $B_\lambda(S)$ is called the *Brandt λ-extension of the semigroup S* [4]. Obviously, if S has zero then $\mathcal{J} = \{0\} \cup \{(\alpha, 0_S, \beta)\colon 0_S \text{ is the zero of } S\}$ is an ideal of $B_\lambda(S)$. We put $B^0_\lambda(S) = B_\lambda(S)/\mathcal{J}$ and the semigroup $B^0_\lambda(S)$ is called the *Brandt λ^0-extension of the semigroup S with zero* [8].

If $\mathcal{I}$ is a trivial semigroup (i.e. $\mathcal{I}$ contains only one element), then we denote the semigroup $\mathcal{I}$ with the adjoined zero by $\mathcal{I}^0$. Obviously, for any $\lambda \geqslant 2$, the Brandt λ^0-extension of the semigroup $\mathcal{I}^0$ is isomorphic to the semigroup of $\lambda\times\lambda$-matrix units and any Brandt λ^0-extension of a semigroup with zero which also contains a non-zero idempotent contains the semigroup of $\lambda\times\lambda$-matrix units. We shall denote the semigroup of $\lambda\times\lambda$-matrix units by B_λ. The 2×2-matrix semigroup with adjoined identity B_2^1 plays an impotent role in Graph Theory and its called the *Perkins semigroup*. In the paper [20] Perkins showed that the semigroup B_2^1 is not finitely based. More details on the word problem of the Perkins semigroup via different graphs may be found in the works of Kitaev and his coauthors (see [17, 18]).

Date: September 21, 2016.

2010 *Mathematics Subject Classification.* 20M15, 20F29 .

Key words and phrases. Semigroup, group of automorphisms, monoid, extension.

We always consider the Brandt λ^0-extension only of a monoid with zero. Obviously, for any monoid S with zero we have $B_1^0(S) = S$. Note that every Brandt λ-extension of a group G is isomorphic to the Brandt λ^0-extension of the group G^0 with adjoined zero. The Brandt λ^0-extension of the group with adjoined zero is called a *Brandt semigroup* [2, 21]. A semigroup S is a Brandt semigroup if and only if S is a completely 0-simple inverse semigroup [1, 19] (cf. also [21, Theorem II.3.5]). We shall say that the Brandt λ^0-extension $B_\lambda^0(S)$ of a semigroup S is *finite* if the cardinal λ is finite.

In the paper [14] Gutik and Repovš established homomorphisms of the Brandt λ^0-extensions of monoids with zeros. They also described a category whose objects are ingredients in the constructions of the Brandt λ^0-extensions of monoids with zeros. Here they introduced finite, compact topological Brandt λ^0-extensions of topological semigroups and countably compact topological Brandt λ^0-extensions of topological inverse semigroups in the class of topological inverse semigroups, and established the structure of such extensions and non-trivial continuous homomorphisms between such topological Brandt λ^0-extensions of topological monoids with zero. There they also described a category whose objects are ingredients in the constructions of finite (compact, countably compact) topological Brandt λ^0-extensions of topological monoids with zeros. These investigations were continued in [10] and [9], where established countably compact topological Brandt λ^0-extensions of topological monoids with zeros and pseudocompact topological Brandt λ^0-extensions of semitopological monoids with zeros their corresponding categories. Some other topological aspects of topologizations, embeddings and completions of the semigroup of $\lambda\times\lambda$-matrix units and Brandt λ^0-extensions as semitopological and topological semigroups were studied in [3, 5, 7, 11, 12, 13, 15, 16].

In this paper we describe the group of automorphisms of the Brandt λ^0-extension $B_\lambda^0(S)$ of an arbitrary monoid S with zero.

2. Automorphisms of the Brandt λ^0-extension of a monoid with zero

We observe that if $f\colon S \to S$ is an automorphism of the semigroup S without zero then it is obvious that the map $\widehat{f}\colon S^0 \to S^0$ defined by the formula

$$(s)\widehat{f} = \begin{cases} (s)f, & \text{if } s \neq 0_S; \\ 0_S, & \text{if } s = 0_S, \end{cases}$$

is an automorphism of the semigroup S^0 with adjoined zero 0_S. Also the automorphism $f\colon S \to S$ of the semigroup S can be extended to an automorphism $f_B\colon B_\lambda^0(S) \to B_\lambda^0(S)$ of the Brandt λ^0-extension $B_\lambda^0(S)$ of the semigroup S by the formulae:

$$(\alpha, s, \beta)f_B = (\alpha, (s)f, \beta), \qquad \text{for all } \alpha, \beta \in \lambda$$

and $(0)f_B = 0$. We remark that so determined extended automorphism is not unique.

The following theorem describes all automorphisms of the Brandt λ^0-extension $B_\lambda^0(S)$ of a monoid S.

Theorem 1. *Let $\lambda \geqslant 1$ be cardinal and let $B_\lambda^0(S)$ be the Brandt λ^0-extension of monoid S with zero. Let $h\colon S \to S$ be an automorphism and suppose that $\varphi\colon \lambda \to \lambda$ is a bijective map. Let H_1 be the group of units of S and $u\colon \lambda \to H_1$ a map. Then the map $\sigma\colon B_\lambda^0(S) \to B_\lambda^0(S)$ defined by the formulae*

$$((\alpha, s, \beta))\sigma = ((\alpha)\varphi, (\alpha)u \cdot (s)h \cdot ((\beta)u)^{-1}, (\beta)\varphi) \qquad \text{and} \qquad (0)\sigma = 0, \tag{1}$$

is an automorphism of the semigroup $B_\lambda^0(S)$. Moreover, every automorphism of $B_\lambda^0(S)$ can be constructed in this manner.

Proof. A simple verification shows that σ is an automorphism of the semigroup $B_\lambda^0(S)$.

Let $\sigma\colon B_\lambda^0(S) \to B_\lambda^0(S)$ be an isomorphism. We fix an arbitrary $\alpha \in \lambda$.

Since $\sigma\colon B_\lambda^0(S) \to B_\lambda^0(S)$ is the automorphism and the idempotent $(\alpha, 1_S, \alpha)$ is maximal with the respect to the natural partial order on $E(B_\lambda^0(S))$, Proposition 3.2 of [14] implies that $((\alpha, 1_S, \alpha))\sigma = (\alpha', 1_S, \alpha')$ for some $\alpha' \in \lambda$.

Since $(\beta, 1_S, \alpha)(\alpha, 1_S, \alpha) = (\beta, 1_S, \alpha)$ for any $\beta \in \lambda$, we have that

$$((\beta, 1_S, \alpha))\sigma = ((\beta, 1_S, \alpha))\sigma \cdot (\alpha', 1_S, \alpha'),$$

and hence

$$((\beta, 1_S, \alpha))\sigma = ((\beta)\varphi, (\beta)u, \alpha'),$$

for some $(\beta)\varphi \in \lambda$ and $(\beta)u \in S$. Similarly, we get that

$$((\alpha, 1_S, \beta))\sigma = (\alpha', (\beta)v, (\beta)\psi),$$

for some $(\beta)\psi \in \lambda$ and $(\beta)v \in S$. Since $(\alpha, 1_S, \beta)(\beta, 1_S, \alpha) = (\alpha, 1_S, \alpha)$, we have that

$$(\alpha', 1_S, \alpha') = ((\alpha, 1_S, \alpha))\sigma = (\alpha', (\beta)v, (\beta)\psi) \cdot ((\beta)\varphi, (\beta)u, \alpha') = (\alpha', (\beta)v \cdot (\beta)u, \alpha'),$$

and hence $(\beta)\varphi = (\beta)\psi = \beta' \in \lambda$ and $(\beta)v \cdot (\beta)u = 1_S$. Similarly, since $(\beta, 1_S, \alpha) \cdot (\alpha, 1_S, \beta) = (\beta, 1_S, \beta)$, we see that the element

$$((\beta, 1_S, \beta))\sigma = ((\beta, 1_S, \alpha)(\alpha, 1_S, \beta))\sigma = (\beta', (\beta)v \cdot (\beta)u, \beta')$$

is a maximal idempotent of the subsemigroup $S_{\beta',\beta'}$ of $B_\lambda^0(S)$, and hence we have that $(\beta)v \cdot (\beta)u = 1_S$. This implies that the elements $(\beta)v$ and $(\beta)u$ are mutually invertible in H_1, and hence $(\beta)v = ((\beta)u)^{-1}$.

If $(\gamma)\varphi = (\delta)\varphi$ for $\gamma, \delta \in \lambda$ then

$$0 \neq (\alpha', 1_S, (\gamma)\varphi) \cdot ((\delta)\varphi, 1_S, \alpha') = ((\alpha, 1_S, \gamma))\sigma \cdot ((\delta, 1_S, \alpha))\sigma,$$

and since σ is an automorphism, we have that

$$(\alpha, 1_S, \gamma) \cdot (\delta, 1_S, \alpha) \neq 0$$

and hence $\gamma = \delta$. Thus $\varphi\colon \lambda \to \lambda$ is a bijective map.

Therefore for $s \in S \setminus \{0_S\}$ we have

$$\begin{aligned}((\gamma, s, \delta))\sigma &= ((\gamma, 1_S, \alpha) \cdot (\alpha, s, \alpha) \cdot (\alpha, 1_S, \delta))\sigma = \\ &= ((\gamma, 1_S, \alpha))\sigma \cdot ((\alpha, s, \alpha))\sigma \cdot ((\alpha, 1_S, \delta))\sigma = \\ &= ((\gamma)\varphi, (\gamma)u, \alpha') \cdot (\alpha', (s)h, \alpha') \cdot (\alpha', ((\delta)u)^{-1}, (\delta)\varphi) = \\ &= ((\gamma)\varphi, (\gamma)u \cdot (s)h \cdot ((\delta)u)^{-1}, (\delta)\varphi).\end{aligned}$$

Also, since 0 is zero of the semigroup $B_\lambda^0(S)$ we conclude that $(0)\sigma = 0$. □

Theorem 1 implies the following corollary:

Corollary 1. *Let $\lambda \geqslant 1$ be cardinal and let $B_\lambda(G)$ be the Brandt semigroup. Let $h\colon G \to G$ be an automorphism and suppose that $\varphi\colon \lambda \to \lambda$ is a bijective map. Let $u\colon \lambda \to G$ be a map. Then the map $\sigma\colon B_\lambda(G) \to B_\lambda(G)$ defined by the formulae*

$$((\alpha, s, \beta))\sigma = ((\alpha)\varphi, (\alpha)u \cdot (s)h \cdot ((\beta)u)^{-1}, (\beta)\varphi) \qquad \text{and} \qquad (0)\sigma = 0,$$

is an automorphism of the Brandt semigroup $B_\lambda(G)$. Moreover, every automorphism of $B_\lambda(G)$ can be constructed in this manner.

Also, we observe that Corollary 1 implies the following well known statement:

Corollary 2. *Let $\lambda \geqslant 1$ be cardinal and $\varphi\colon \lambda \to \lambda$ a bijective map. Then the map $\sigma\colon B_\lambda \to B_\lambda$ defined by the formulae*

$$((\alpha, \beta))\sigma = ((\alpha)\varphi, (\beta)\varphi) \quad \text{and} \quad (0)\sigma = 0,$$

is an automorphism of the semigroup of $\lambda\times\lambda$-matrix units B_λ. Moreover, every automorphism of B_λ can be constructed in this manner.

The following example implies that the condition that semigroup S contains the identity is essential.

Example 1. Let λ be any cardinal $\geqslant 2$. Let S be the zero-semigroup of cardinality $\geqslant 3$ and 0_S is zero of S. It is easily to see that every bijective map $\sigma\colon B_\lambda^0(S) \to B_\lambda^0(S)$ such that $(0)\sigma = 0$ is an automorphism of the Brandt λ^0-extension of S.

Remark. By Theorem 1 we have that every automorphism $\sigma\colon B_\lambda^0(S) \to B_\lambda^0(S)$ of the Brandt λ^0-extension of an arbitrary monoid S with zero identifies with the ordered triple $[\varphi, h, u]$, where $h\colon S \to S$ is an automorphism of S, $\varphi\colon \lambda \to \lambda$ is a bijective map and $u\colon \lambda \to H_1$ is a map, where H_1 is the group of units of S.

Lemma 1. *Let $\lambda \geqslant 1$ be cardinal, S be a monoid with zero and let $B_\lambda^0(S)$ be the Brandt λ^0-extension of S. Then the composition of arbitrary automorphisms $\sigma = [\varphi, h, u]$ and $\sigma' = [\varphi', h', u']$ of the Brandt λ^0-extension of S defines in the following way:*

$$[\varphi, h, u] \cdot [\varphi', h', u'] = [\varphi\varphi', hh', \varphi u' \cdot uh'].$$

Proof. By Theorem 1 for every $(\alpha, s, \beta) \in B_\lambda^0(S)$ we have that

$$\begin{aligned}(\alpha, s, \beta)(\sigma\sigma') &= \big((\alpha)\varphi, (\alpha)u\cdot(s)h\cdot((\beta)u)^{-1}, (\beta)\varphi\big)\,\sigma' = \\ &= \big(((\alpha)\varphi)\varphi', ((\alpha)\varphi)u' \cdot \big((\alpha)u \cdot (s)h \cdot ((\beta)u)^{-1}\big)\, h' \cdot (((\beta)\varphi)u')^{-1}, ((\beta)\varphi)\varphi'\big) =\end{aligned}$$

and since h' is an automorphism of the monoid S we get that this is equal to

$$\begin{aligned}&= \big(((\alpha)\varphi)\varphi', ((\alpha)\varphi)u' \cdot ((\alpha)u)\, h' \cdot ((s)h)\, h' \cdot (((\beta)u)h')^{-1} \cdot (((\beta)\varphi)u')^{-1}, ((\beta)\varphi)\varphi'\big) = \\ &= \big((\alpha)(\varphi\varphi'), (\alpha)(\varphi u' \cdot uh') \cdot ((s)h)\, h' \cdot (\beta)\,(\varphi u' \cdot uh')^{-1}, (\beta)(\varphi\varphi')\big).\end{aligned}$$

This completes the proof of the requested equality. □

Theorem 2. *Let $\lambda \geqslant 1$ be cardinal, S be a monoid with zero and let $B_\lambda^0(S)$ be the Brandt λ^0-extension of S. Then the group of automorphisms $\mathbf{Aut}(B_\lambda^0(S))$ of $B_\lambda^0(S)$ is isomorphic to a homomorphic image of the group defines on the Cartesian product $\mathscr{S}_\lambda \times \mathbf{Aut}(S) \times H_1^\lambda$ with the following binary operation:*

$$[\varphi, h, u] \cdot [\varphi', h', u'] = [\varphi\varphi', hh', \varphi u' \cdot uh'], \tag{2}$$

where $\mathscr{S}_\lambda$ is the group of all bijections of the cardinal λ, $\mathbf{Aut}(S)$ is the group of all automorphisms of the semigroup S and H_1^λ is the direct λ-power of the group of units H_1 of the monoid S. Moreover, the inverse element of $[\varphi, h, u]$ in the group $\mathbf{Aut}(B_\lambda^0(S))$ is defined by the formula:

$$[\varphi, h, u]^{-1} = \left[\varphi^{-1}, h^{-1}, \varphi^{-1}u^{-1}h^{-1}\right].$$

Proof. First, we show that the binary operation defined by formula (2) is associative. Let $[\varphi, h, u]$, $[\varphi', h', u']$ and $[\varphi'', h'', u'']$ be arbitrary elements of the Cartesian product $\mathscr{S}_\lambda \times \mathbf{Aut}(S) \times H_1^\lambda$. Then we have that

$$\begin{aligned}\big([\varphi, h, u] \cdot [\varphi', h', u']\big) \cdot [\varphi'', h'', u''] &= [\varphi\varphi', hh', \varphi u' \cdot uh'] \cdot [\varphi'', h'', u''] = \\ &= [\varphi\varphi'\varphi'', hh'h'', \varphi\varphi' u'' \cdot (\varphi u' \cdot uh')h''] = \\ &= [\varphi\varphi'\varphi'', hh'h'', \varphi\varphi' u'' \cdot \varphi u'h'' \cdot uh'h'']\end{aligned}$$

and

$$\begin{aligned}[\varphi, h, u] \cdot ([\varphi', h', u'] \cdot [\varphi'', h'', u'']) &= [\varphi, h, u] \cdot [\varphi'\varphi'', h'h'', \varphi' u'' \cdot u'h''] = \\ &= [\varphi\varphi'\varphi'', hh'h'', \varphi(\varphi' u'' \cdot u'h'') \cdot uh'h''] = \\ &= [\varphi\varphi'\varphi'', hh'h'', \varphi\varphi' u'' \cdot \varphi u'h'' \cdot uh'h''],\end{aligned}$$

and hence so defined operation is associative.

Theorem 1 implies that formula (1) determines a map $\mathfrak{F}$ from the Cartesian product $\mathscr{S}_\lambda \times \mathbf{Aut}(S) \times H_1^\lambda$ onto the group of automorphisms $\mathbf{Aut}(B_\lambda^0(S))$ of the Brandt λ^0-extension $B_\lambda^0(S)$ of the monoid S, and hence the associativity of binary operation (2) implies that the map $\mathfrak{F}$ is a homomorphism from $\mathscr{S}_\lambda \times \mathbf{Aut}(S) \times H_1^\lambda$ onto the group $\mathbf{Aut}(B_\lambda^0(S))$.

Next we show that $[1_{\mathscr{S}_\lambda}, 1_{\mathbf{Aut}(S)}, 1_{H_1^\lambda}]$ is a unit element with the respect to the binary operation (2), where $1_{\mathscr{S}_\lambda}$, $1_{\mathbf{Aut}(S)}$ and $1_{H_1^\lambda}$ are units of the groups $\mathscr{S}_\lambda$, $\mathbf{Aut}(S)$ and H_1^λ, respectively. Then we have

that

$$\begin{aligned}[\varphi,h,u]\cdot\left[1_{\mathscr{S}_\lambda},1_{\mathbf{Aut}(S)},1_{H_1^\lambda}\right] &= \left[\varphi 1_{\mathscr{S}_\lambda}, h1_{\mathbf{Aut}(S)}, \varphi 1_{H_1^\lambda}\cdot u1_{\mathbf{Aut}(S)}\right] = \\ &= \left[\varphi, h, \varphi 1_{H_1^\lambda}\cdot u1_{\mathbf{Aut}(S)}\right] = \\ &= \left[\varphi, h, 1_{H_1^\lambda}\cdot u\right] = \\ &= [\varphi,h,u]\end{aligned}$$

and

$$\left[1_{\mathscr{S}_\lambda},1_{\mathbf{Aut}(S)},1_{H_1^\lambda}\right]\cdot[\varphi,h,u] = \left[1_{\mathscr{S}_\lambda}\varphi, 1_{\mathbf{Aut}(S)}h, 1_{\mathscr{S}_\lambda}u\cdot 1_{H_1^\lambda}h\right] = [\varphi,h,u],$$

because every automorphism $h\in\mathbf{Aut}(S)$ acts on the group H_1^λ by the natural way as a restriction of global automorphism of the semigroup S on every factor, and hence we get that $1_{H_1^\lambda}h = 1_{H_1^\lambda}$.

Also, similar arguments imply that

$$\begin{aligned}[\varphi,h,u]\cdot[\varphi,h,u]^{-1} &= [\varphi,h,u]\cdot\left[\varphi^{-1},h^{-1},\varphi^{-1}u^{-1}h^{-1}\right] = \\ &= \left[\varphi\varphi^{-1}, hh^{-1}, (\varphi\varphi^{-1})u^{-1}h^{-1}\cdot uh^{-1}\right] = \\ &= \left[\varphi\varphi^{-1}, hh^{-1}, (1_{\mathscr{S}_\lambda})u^{-1}h^{-1}\cdot uh^{-1}\right] = \\ &= \left[\varphi\varphi^{-1}, hh^{-1}, u^{-1}h^{-1}\cdot uh^{-1}\right] = \\ &= \left[1_{\mathscr{S}_\lambda},1_{\mathbf{Aut}(S)},1_{H_1^\lambda}\right]\end{aligned}$$

and

$$\begin{aligned}[\varphi,h,u]^{-1}\cdot[\varphi,h,u] &= \left[\varphi^{-1},h^{-1},\varphi^{-1}u^{-1}h^{-1}\right]\cdot[\varphi,h,u] = \\ &= \left[\varphi^{-1}\varphi, h^{-1}h, \varphi^{-1}u\cdot\varphi^{-1}u^{-1}h^{-1}h\right] = \\ &= \left[\varphi^{-1}\varphi, h^{-1}h, \varphi^{-1}u\cdot\varphi^{-1}u^{-1}\right] = \\ &= \left[1_{\mathscr{S}_\lambda},1_{\mathbf{Aut}(S)},1_{H_1^\lambda}\right].\end{aligned}$$

This implies that the elements $[\varphi^{-1},h^{-1},\varphi^{-1}u^{-1}h^{-1}]$ and $[\varphi,h,u]$ are invertible in $\mathscr{S}_\lambda\times\mathbf{Aut}(S)\times H_1^\lambda$, and hence the set $\mathscr{S}_\lambda\times\mathbf{Aut}(S)\times H_1^\lambda$ with the binary operation (2) is a group.

Let $\mathbf{Id}\colon B_\lambda^0(S)\to B_\lambda^0(S)$ be the identity automorphism of the semigroup $B_\lambda^0(S)$. Then by Theorem 1 there exist some automorphism $h\colon S\to S$, a bijective map $\varphi\colon\lambda\to\lambda$ and a map $u\colon\lambda\to H_1$ into the group H_1 of units of S such that

$$(\alpha,s,\beta) = (\alpha,s,\beta)\mathbf{Id} = ((\alpha)\varphi, (\alpha)u\cdot(s)h\cdot((\beta)u)^{-1}, (\beta)\varphi),$$

for all $\alpha,\beta\in\lambda$ and $s\in S^*$. Since $\mathbf{Id}\colon B_\lambda^0(S)\to B_\lambda^0(S)$ is the identity automorphism we conclude that $(\alpha)\varphi=\alpha$ for every $\alpha\in\lambda$. Also, for every $s\in S^*$ we get that $s=(\alpha)u\cdot(s)h\cdot((\beta)u)^{-1}$ for all $\alpha,\beta\in\lambda$, and hence we obtain that

$$1_S = (\alpha)u\cdot(1_S)h\cdot((\beta)u)^{-1} = (\alpha)u\cdot((\beta)u)^{-1}$$

for all $\alpha,\beta\in\lambda$. This implies that $(\alpha)u=(\beta)u=\widetilde{u}$ is a fixed element of the group H_1 for all $\alpha,\beta\in\lambda$.

We define

$$\ker N = \left\{[\varphi,h,\widetilde{u}]\in\mathscr{S}_\lambda\times\mathbf{Aut}(S)\times H_1^\lambda\colon \varphi\colon\lambda\to\lambda \text{ is an idemtity map, } \widetilde{u}(s)h\widetilde{u}^{\,-1}=s \text{ for any } s\in S\right\}.$$

It is obvious that the equality $\widetilde{u}(s)h\widetilde{u}^{-1}=s$ implies that $(s)h=\widetilde{u}^{-1}s\widetilde{u}$ for all $s\in S$. The previous arguments implies that $[\varphi,h,\widetilde{u}]\in\ker N$ if and only if $[\varphi,h,\widetilde{u}]\mathfrak{F}$ is the unit of the group $\mathbf{Aut}(B_\lambda^0(S))$, and hence $\ker N$ is a normal subgroup of $\mathscr{S}_\lambda\times\mathbf{Aut}(S)\times H_1^\lambda$. This implies that the quotient group $(\mathscr{S}_\lambda\times\mathbf{Aut}(S)\times H_1^\lambda)/\ker N$ is isomorphic to the group $\mathbf{Aut}(B_\lambda^0(S))$. □

References

[1] A. H. Clifford, *Matrix representations of completely simple semigroups*, Amer. J. Math. **64** (1942), 327–342.
[2] A. H. Clifford and G. B. Preston, *The Algebraic Theory of Semigroups*, Vols. I and II, Amer. Math. Soc. Surveys **7**, Providence, R.I., 1961 and 1967.
[3] S. Bardyla and O. Gutik, *On a semitopological polycyclic monoid*, Algebra Discr. Math. **21**:2 (2016), 163–183.
[4] O. V. Gutik, *On Howie semigroup*, Mat. Metody Phis.-Mekh. Polya. **42**:4 (1999), 127–132 (in Ukrainian).
[5] O. Gutik, *On closures in semitopological inverse semigroups with continuous inversion*, Algebra Discr. Math. **18**:1 (2014), 59–85.
[6] O. V. Gutik and K. P. Pavlyk, *Topological Brandt λ-extensions of absolutely H-closed topological inverse semigroups*, Visnyk Lviv Univ., Ser. Mekh.-Math. **61** (2003), 98–105.
[7] O. V. Gutik and K. P. Pavlyk, *On topological semigroups of matrix units*, Semigroup Forum **71**:3 (2005), 389–400.
[8] O. V. Gutik and K. P. Pavlyk, *On Brandt λ^0-extensions of semigroups with zero*, Mat. Metody Phis.-Mekh. Polya. **49**:3 (2006), 26–40.
[9] O. Gutik and K. Pavlyk, *On pseudocompact topological Brandt λ^0-extensions of semitopological monoids*, Topological Algebra Appl. **1** (2013), 60–79.
[10] O. Gutik, K. Pavlyk, and A. Reiter, *Topological semigroups of matrix units and countably compact Brandt λ^0-extensions*, Mat. Stud. **32**:2 (2009), 115–131.
[11] O. V. Gutik, K. P. Pavlyk, and A. R. Reiter, *On topological Brandt semigroups*, Mat. Metody Fiz.-Mekh. Polya **54**:2 (2011), 7–16 (in Ukrainian); English version in: J. Math. Sci. **184**:1 (2012), 1–11.
[12] O. Gutik and O. Ravsky, *On feebly compact inverse primitive (semi)topological semigroups*, Mat. Stud. **44**:1 (2015), 3–26.
[13] O. V. Gutik and O. V. Ravsky, *Pseudocompactness, products and Brandt λ^0-extensions of semitopological monoids*, Mat. Metody Fiz.-Mekh. Polya **58**:2 (2015), 20–37.
[14] O. Gutik and D. Repovš, *On Brandt λ^0-extensions of monoids with zero*, Semigroup Forum **80**:1 (2010), 8–32.
[15] J. Jamalzadeh and Gh. Rezaei, *Countably compact topological semigroups versus Brandt extensions and paragroups*, Algebras Groups Geom. **27**:2 (2010), 219–228.
[16] J. Jamalzadeh and Gh. Rezaei, *Brandt extensions and primitive topologically periodic inverse topological semigroups*, Bull. Iran. Math. Soc. **39**:1 (2013), 87–95.
[17] S. Kitaev and V. Lozin, *Words and Graphs*, Monographs in Theor. Comput. Sc. An EATCS Series. Springer, Cham, 2015.
[18] S. Kitaev and S. Seif, *Word problem of the Perkins semigroup via directed acyclic graphs*, Order **25**:3 (2008), 177–194.
[19] W. D. Munn, *Matrix representations of semigroups*, Proc. Cambridge Phil. Soc. **53** (1957), 5–12.
[20] P. Perkins, *Bases for equational theories of semigroups*, J. Algebra **11**:2 (1969), 298–314.
[21] M. Petrich, *Inverse Semigroups*, John Wiley & Sons, New York, 1984.

Faculty of Mathematics, National University of Lviv, Universytetska 1, Lviv, 79000, Ukraine
E-mail address: o_gutik@franko.lviv.ua, ovgutik@yahoo.com